



## **NH Winter Climate and the AO**

Forecasting the phase of the Arctic Oscillation is a factor for seasonal and longer-range forecasts.





**Future climate change** 

- (1) Greenhouse gas forcing predicts more +AO conditions (e.g., Shindell et al. 1999, Hurrell et al. 2004; Miller et al. 2006, Scaife et al. 2012).
- (2) Will winters grow increasingly warmer? Or are there important feedbacks that could mitigate the positive trend, even intermittently?



## **Fall Eurasian Snow-Winter AO Hypothesis**

October Siberian Snow Cover Expands SLP Increases Over Eurasia November Vertical Wave Propagation (i.e., Poleward Heat Fluxes) Enhanced December Stratospheric Polar Vortex Weakens or Breaks Down Stratospheric Circulation Ianuary **Anomalies Propagate Downward** into the Troposphere **Negative AO Conditions** February Prevail in the Troposphere Based on Cohen et al. (2007)

Evidenced in observations (e.g., Foster et al. 1983; Cohen and Entekhabu 1998; Saito et al. 2001; Cohen et al. 2007).

Model-produced snow cover does *not* demonstrate the response (e.g., *Hardiman et al.* 2008; *Allen and Zender* 2011), but a model with prescribed snow can (e.g., *Fletcher et al.* 2009; *Allen and Zender* 2010, 2011).

#### **Objectives**

- 1) Examine salient features for the mechanism in the models (e.g., snow cover, AO).
- 2) Evaluate the 'six-step process' in CMIP5 models and compare with observations.
- 3) Offer suggestions for why CMIP5 models do not agree with observations.



## **Data and Methodology**

#### **Observational Data**


- •Monthly-mean ERA-Interim (1979-2011)
- •October Monthly-Mean Rutgers Eurasian Snow Cover Index (20-75°N, 0-170°E) (1979-2010)

#### **CMIP5 Models**

- •Monthly-mean piControl runs (15 models).
- •Selected based on availability of snow cover extent (snc) as downloadable variable.
- •Regridded to a 2.5° by 2.5° grid for intermodel comparisons.








#### Methodology

- Subdivide the piControl runs into 40-yr segments.
- Compute statistics on each segment separately.
- Present results by model (aggregate segment statistics) and as 'multi-model ensemble-mean.'
- Focus on NH extended cold season (ONDJFM).




## The AO Pattern – Obs. vs. CMIP5

## SLPa Regressed on -PC1 Of NDJFM SLPa



# October Eurasian Snow Cover Statistics


#### October Mean Eurasian Snow Cover Extent





#### **October Eurasian Snow Cover Statistics**

#### Standard Deviation - October Eurasian Snow Cover






#### **Eurasian Snow/SLP Relations**

Atmospheric and Environmental Research ND SLPa Regressed Onto Oct. Eurasian Snow Cover Siberian Snow Cover Expands


2

SLP Increases Over Eurasia




## **Eurasian Snow/Surf. T Relations**

ND Surface Ta Regressed Onto Oct. Eurasian Snow Cover




## Steps 3 – 5 of the Hypothesis



## Step 6 - Link to the DJF AO

#### Correlation of DJF AO Index w/ Oct Snow Index






#### (1) Variability in the Stratospheric Polar Vortex


- All models show lower variability than observed, some significantly lower (e.g., CSIRO).
- Ensemble-mean  $\sigma_{U60} = 9.1$  m/s vs. Observed  $\sigma_{U60} = 15$  m/s.

#### Standard Deviation of January [U]<sub>10</sub> at 60°N







#### (2) Downward Propagation of Stratospheric Anomalies



Ensemble-mean correlations show downward propagation to ~150 hPa but not to the surface.

r(Jan AO<sub>10</sub>, Jan AO<sub>1000</sub>) ranges from -0.54 (INMCM4) to 0.20 in MIROC5.







## **Summary and Conclusions**

- The six-step snow-AO hypothesis does not verify in the CMIP5 models, similar to the results from the CMIP3 models (*Hardiman et al.* 2008).
- Models continue to underperform on simulating fall snow cover extent, its variability, and the lagged atmospheric response to the snow.
- Analysis with the <u>historical</u> runs yields very similar conclusions.
- Irrespective of the snow relationship, the coupled climate models have issues with stratospheric vortex variability and 'downward propagation'.
  - This fact may give pause for wintertime climate model projections.

#### • Remaining Challenges/Future Work

- Snowfall/snow cover in the models. Precipitation-related? Land surface?
- Investigation of daily-mean output for downward propagation and wave dynamics propagation. This is relevant for both S/T studies as well as the snow-AO hypothesis.



## 3 Month Seasonal Forecast: U.S.

# Thank you!

Jason C Furtado

Atmospheric and Environmental Research

E-mail: jfurtado@aer.com

This work is funded through NOAA Grant #NA10OAR4310163 and NSF Grant #BCS-1060323.






#### **EXTRA SLIDES**

October 24, 2012 Jason C. Furtado, Atmospheric and Environmental Research, Inc. (AER) NOAA 37<sup>TH</sup> Annual Climate Diagnostics Workshop



## **Prior Work**


# **Observations** a) Corr Oct Snow and 40-80N WAF 100 b) Corr Oct Snow and 60-90N GPH 1000 Dec Jan Day Cohen et al. (2007) Oct Nov





#### (1) Wave Forcing and Wave Propagation

Oct Zonal-Mean U (shading) and EP-Fluxes (arrows) Regressed onto the Oct Snow Index





#### (2) ND WAFz / JF SLPa Covariability

